

for European patients with advanced EGFR mutation-positive non-small-cell lung cancer (EURTAC): a multicentre, open-label, randomised phase 3 trial. *Lancet Oncol* 2012;13:239–246.

3. Zhou C, Wu YL, Chen G, et al. Erlotinib versus chemotherapy as first-line treatment for patients with advanced EGFR mutation-positive non-small-cell lung cancer (OPTIMAL, CTONG-0802): a multicentre, open-label, randomised, phase 3 study. *Lancet Oncol* 2011;12:735–742.
4. Hata A, Katakami N, Yoshioka H, et al. How sensitive are epidermal growth factor receptor-tyrosine kinase inhibitors for squamous cell carcinoma of the lung harboring EGFR gene-sensitive mutations? *J Thorac Oncol* 2013;8:89–95.
5. Sequist LV, Yang JC, Yamamoto N, et al. Phase III study of afatinib or cisplatin plus pemetrexed in patients with metastatic lung adenocarcinoma with EGFR mutations. *J Clin Oncol* 2013;31:3327–3334.

How Sensitive Are Epidermal Growth Factor Receptor–Tyrosine Kinase Inhibitors for Squamous Cell Carcinoma of the Lung Harboring EGFR Gene–Sensitive Mutations?

In Response:

We are thankful for the thoughtful suggestions by Zhou et al.¹ They raised two intriguing questions: first, which is better, platinum doublets or epidermal growth factor receptor (EGFR)–tyrosine kinase inhibitor (TKI) as first-line treatment in squamous cell carcinoma (SQC) patients with EGFR-sensitive mutations; and second, how do we treat patients with minor EGFR mutations?

Disclosure: The author declares no conflict of interest.

Address for correspondence: Akito Hata, MD, Division of Integrated Oncology, Institute of Biomedical Research and Innovation, 2-2, Minatojima-Minamimachi, Chuo-ku, Kobe, 650-0047, Japan. E-mail: a-hata@fbri.org

Copyright © 2013 by the International Association for the Study of Lung Cancer
ISSN: 1556-0864/14/0902-0e20

Several studies in patients with *EGFR* mutations demonstrated the efficacy of platinum doublets; and the response rate (RR) and median progression-free survival (PFS) were approximately 30% and 6 months,² similar to our 14 *EGFR*-mutated SQC patients receiving platinum doublets as first-line treatment. Compared with our EGFR-TKI results (RR, 25% and median PFS, 1.4 months),³ we would generally choose platinum doublets as first-line treatment before EGFR-TKIs in such patients. However, we should ponder how many *EGFR*-mutated adenocarcinoma cells are included in an SQC case. If the *EGFR*-mutated SQC patient is suspected as harboring *EGFR*-mutated adenocarcinoma propensity per the following factors: female, never smoker, high carcinoembryonic antigen, and thyroid transcription factor-1 positive, we could try EGFR-TKI as a first-line treatment. Conversely, male, current smoker, high cytokeratin 19 fragment, and p63 positive patients would be better served by platinum doublets. We should recognize some *EGFR*-mutated SQC include an incomplete sampling of adeno-SQC and/or poorly differentiated adenocarcinoma morphologically mimicking SQC. Approximately one third of the *EGFR*-mutated SQC patients in our study obtained a PFS of longer than 6 months, and most of them had adenocarcinoma propensities. To better identify EGFR-TKI-sensitive SQC patients, not only *EGFR* mutation status, but also clinical factors and pathologic findings should be taken into consideration.

Unfortunately, both of our two patients with a G719X point mutation in exon 18 did not undergo EGFR-TKIs in our study. Sensitivities of EGFR-TKIs in minor *EGFR* mutations such as G719X seem to be less effective than major mutations such as deletional mutations in exon 19 and a L858R point mutation in exon 21. In patients with minor mutations, the RR and PFS were shown to be 30% to 50% and 3 to 5 months, comparable to platinum doublets.⁴ In our opinion, we would choose platinum doublets as first-line treatment,

before EGFR-TKI in patients with minor mutations. It is more difficult to administer platinum doublets as second-line treatment, compared with EGFR-TKI. In patients with SQC harboring minor mutations, priority of EGFR-TKI becomes much lower.

The diagnostic limitations of small biopsies and intratumoral heterogeneity make definitive diagnoses difficult.⁵ Clinicians should understand the limitations of pathologic diagnosis based on small biopsies, and need to cooperate with pathologists to better reach a diagnosis for individual patients. EGFR-TKI sensitivity in *EGFR*-mutated SQC should be judged by the general factors of the patient.

Akito Hata, MD

Nobuyuki Katakami, MD, PhD

Division of Integrated Oncology
Institute of Biomedical Research and Innovation
Kobe, Japan

Hiroshige Yoshioka, MD

Department of Respiratory Medicine
Kurashiki Central Hospital
Kurashiki, Japan

REFERENCES

1. Zhou F, Chen X, Zhou C. How sensitive are epidermal growth factor receptor–tyrosine kinase inhibitors for squamous cell carcinoma of the lung harboring egfr gene–sensitive mutations? *J Thorac Oncol* 2014;9:e19–e20.
2. Mitsudomi T, Morita S, Yatabe Y, et al.; West Japan Oncology Group. Gefitinib versus cisplatin plus docetaxel in patients with non-small-cell lung cancer harbouring mutations of the epidermal growth factor receptor (WJTOG3405): an open label, randomised phase 3 trial. *Lancet Oncol* 2010;11:121–128.
3. Hata A, Katakami N, Yoshioka H, et al. How sensitive are epidermal growth factor receptor-tyrosine kinase inhibitors for squamous cell carcinoma of the lung harboring EGFR gene-sensitive mutations? *J Thorac Oncol* 2013;8:89–95.
4. De Pas T, Toffalorio F, Manzotti M, et al. Activity of epidermal growth factor receptor-tyrosine kinase inhibitors in patients with non-small cell lung cancer harboring rare epidermal growth factor receptor mutations. *J Thorac Oncol* 2011;6:1895–1901.
5. Travis WD, Rekhtman N, Riley GJ, et al. Pathologic diagnosis of advanced lung cancer based on small biopsies and cytology: a paradigm shift. *J Thorac Oncol* 2010;5:411–414.